

THE MEADOW, PSAMMOPHYTIC AND RUDERAL PLANT COMMUNITIES WITH *SOLIDAGO CANADENSIS* L. IN CHERNIHIV POLESIE (UKRAINE)

**Hanna Danko^{*}, Oleksandr Lukash, Iryna Morozova, Volodymyr Boiko,
Oleksandr Yakovenko**

T.H. Shevchenko National University “Chernihiv Colegium” Hetman Polubotok Str. 53, 14013 Chernihiv, Ukraine;
e-mails: H. Danko <annadanko978@gmail.com>, O. Lukash <lukash2011@ukr.net>, V. Boiko <boiko_95@ukr.net>,
O. Yakovenko <ajakov2@gmail.com>, I. Morozova <irina.morozova1102@gmail.com>

* corresponding author



Abstract:

In Chernihiv Polesie *Solidago canadensis* most often grows in ruderal communities of the *Berteroëtum incanae* association. Characteristic plant species of the *Artemisietea vulgaris* class have been found in many phytocenoses with *Solidago canadensis*. A typical ruderal community dominated by *S. canadensis* was found, in which characteristic species of the xero-mesophytic ruderal vegetation of the *Onopordion acanthii* are well represented. Initial communities with the *S. canadensis* coverage of 25 to 60% in combination with the species of this order and the characteristic species of other high syntaxa were found. Most of them are the transformed meadow phytocenoses of the river floodplains and less often – the psammophytic phytocenoses of pine terraces. The process of ruderalization of meadow ecosystems as a result of the invasion of *S. canadensis* in Chernihiv Polesie was revealed. This process is especially pronounced on the loess islands, where meadows change into semiruderal grasslands and herblands of the *Convolvulo arvensis–Agropyron repens* alliance. *S. canadensis* invasion leads to xerophytization and unification of the floodplains meadow phytocenoses grassland. The course of these processes is accelerated by anthropogenic pressure on ecosystems and has irreversible consequences. *S. canadensis* rarely occurs in the *Koelerio–Corynephoretea canescens* class psammophytic communities.

Key words: vegetation, anthropogenic influence, Chernihiv Polesie, *Solidago canadensis* L.

Manuscript received 24 November 2020, accepted 19 July 2021

INTRODUCTION

The extent of ecosystem modification by invasive plants depends on the species and location, hence invasions of the same species may have different effects, depending on local conditions (Dassonville *et al.*, 2008). *Solidago canadensis* is a violent species with significant phenotypic and phenological variability, propagated by achenes and restored by rhizomes. Due to this, it quickly colonizes disturbed areas and successfully masters various types of habitats (Burda *et al.*, 2015). *S. canadensis* has an allelopathic effect on the co-occurring flora and vegetation. This is demonstrated by publications of the last years (Abhilasha *et al.*, 2008). From the other hand *S. canadensis* is an important source of honey during late summer and early autumn (Stefanic *et al.*, 2003).

In the southeast of Belarus *S. canadensis* invaded the plant communities of abandoned agricultural lands at the

primary stages of progressive succession, and can cause its inhibition (Gusev, 2015). The *S. canadensis* communities can break regenerative plant succession in the sands of technogenic landscapes (Gusev and Shpileuskaya, 2016). Despite the slower *S. canadensis* invasion into natural plant communities, this process is more dangerous. It can lead to the inhibition and even complete extinction of native species in protected areas (Dubovik *et al.*, 2019). It was found that presence of *S. canadensis* at two localities in southern (Olkusz) and eastern Poland (Siedlce) reduces a number of vascular plant species and an increase in *S. canadensis* cover results in soil degradation and habitat homogenization (Bielecka *et al.*, 2020).

The study of the syntaxonomic composition of the *S. canadensis* communities is important for assessing its invasive activity in the region under consideration. Analysis of the floristic composition of geobotanical relevés with the dominance of *S. canadensis* performed in the

cities of Kursk, Bryansk and Minsk revealed that they are grouped into the *Solidaginetum serotino-canadensis* association (the *Dauco-Melilotion* alliance), the *Rudbeckio-laciniatae-Solidaginetum canadensis* association (the *Dauco-Melilotion* alliance) and the *Solidago canadensis* initial community (the *Galio-Urticetea* class) (Arepieva and Kulikova, 2017). These groups depend both on humidity and richness of soils and substrates where the communities develop, and the degree of anthropogenic influence on them. In southern Belarus this species forms the *Solidago canadensis* initial community, phytocenoses of the *Calamagrostio epigeiosi-Solidaginetum canadensis* association (the *Dauco-Melilotion* alliance) and it is also a part of the *Arctio-Artemisietum vulgaris* ruderal communities (Daineka and Timofeev, 2018).

S. canadensis and its communities pose a threat to the diversity of the Chernihiv Polesie natural habitats. Our aim was to establish the syntaxonomic composition of plant communities with *S. canadensis* in this region.

REGIONAL SETTING

According to the physical and geographical zoning of Ukraine (Marynich *et al.*, 2003), the northwestern part of the Chernihiv region is situated within the boundaries of Chernihiv Polesie. In the geostructural relation, the Chernihiv Polesie is located within the limits of three structural formations: the eastern slope and the foot of the Ukrainian crystalline shield, the western slope of the Voronezh crystalline massif and the Dnipro-Donets cavity located between them.

Modern landscapes of the Chernihiv Polesie were formed in the Middle and Late Quaternary. The landscape structure of this territory is dominated by the Polesie landscape complexes (63% of the region area). The landscape complexes with forest-steppe features have a significant (up to 18%) distribution. They occur fragmentarily and are characteristic for the so-called 'loess islands'. These landscape complexes spread in a strip located to the north of the Desna River (Lukash *et al.*, 2018).

The landscape features of the Chernihiv Polesie and anthropogenic influence are reflected in the current state of the regional vegetation. The forest and the meadow phytocenoses prevail in the vegetation cover of the region. The psammophytic vegetation is represented fragmentarily. It is possible to observe the adventitization and apophytization of sand vegetation, one of the indicators of which is the rupture of cenotic connections between the species of the community (Lukash and Danko, 2020). The natural vegetation of the loess islands of the Chernihiv Polesie has been largely transformed due to the agricultural use and the vegetal plant communities predominate. Long-term use of the arable lands led not only to the stabilization of the composition of weed synusiae of agro-phytocenoses, but also contributed to the increase of erosion and suffusion forms of the relief (Lukash *et al.*, 2018). The region has an extensive hydrological network – the Dnipro River basins

and its tributaries, the Desna River. Natural vegetation covers about 70% of the total river basins area.

MATERIAL AND METHODS

The materials for the article were collected during the field research of the Chernihiv Polesie vegetation in 2018–2020. The field study of the vegetation was carried out by geobotanical methods (Lavrenko and Korchagin, 1976). The vegetation descriptions were carried out during the optimum of the vegetation period in the areas of 30–100 m² each. Cover abundance scale is the following: + – up to 1%, 1 – 1–5%, 2 – 6–15, 3 – 16–25%, 4 – 26–50%, 5 >50%. The projective cover of *S. canadensis* is given as a percentage. 29 phytosociological relevés were taken. Syntaxa were identified according to Mucina *et al.* (2016), Matuszkiewicz (2019) (for natural vegetation), Solomakha *et al.* (1992) (for synanthropic vegetation). Syntaxon names are ordered according to Mucina *et al.* (2016). The nomenclature of the plant taxa follows the Plant List (www.theplantlist.org). The successional stages of vegetation are named after the dominant species. Geobotanical data were added to a database by means of TURBOVEG 2.79 software (Hennekens, Schamine, 2001), the interpretation of geobotanical material was carried out with use of the software package JUICE 7.0.83 (Tiche, 2002).

RESULTS AND DISCUSSION

According to the results of the field research, which took place in the Chernihiv Polesie (Figs 1 and 2), the syntaxonomic affiliation of the phytocenoses, in which *S. canadensis* grows, has been established (Table 1).

The relevés 1–6 belong to the *Berteroëtum incanae* association of the *Onopordion acanthii* alliance of the *Onopordetalia acanthii* order of the *Artemisietea vulgaris* class. The cenoses of the association occupy the largest areas on the pine-forest terraces on well-drained sandy soils of the Snov and the Stryzhen rivers. The communities are formed by such species as *Berteroëa incana*, *Plantago indica* and sometimes *Centaurea diffusa*. The characteristic species of the *Artemisietea vulgaris* class are also represented. The species of the *Papaveretea rhoeadis* class (*Chenopodium album*, *Erigeron annuus*, *Erigeron canadensis*, *Lepidium densiflorum*, *Setaria pumila*, *Vicia sativa*) accompany this community. In western Europe the *Berteroëtum incanae* association occurs in contact with the *Arrhenatheretalia* order communities, whereas in the eastern Europe it is more frequently found in contact with the *Festuco-Brometea* and *Sedo-Scleranthetea* class (Mucina and Brandes, 1985).

However, the structure and (or) composition of the described communities do not allow to include clearly each of them to one or another association of the *Onopordion acanthii* alliance (thistle-dominated xero-mesophytic ruderal vegetation of subcontinental central Europe and the northern Balkans). It should be noted that species of the




Fig. 1. Location of the investigated plant communities with *Solidago canadensis* participation.



Fig. 2. Location of the investigated plant communities with *Solidago canadensis* participation in Chernihiv.

Artemisietea vulgaris, the *Molinio-Arrhenatheretea*, the *Koelerio-Corynephoretea canescens* and the *Epilobietea angustifoli* classes were diagnosed in the relevés 7–19. The described phytocenoses are probably the communities at early stages of succession with the domination of *S. canadensis*.

One of these communities (rélevé 7) is formed by *Solidago canadensis*, which is monodominant. This community is also characterized by a high participation of biennial or perennial plants, which are characteristic species of the *Onopordetalia acanthii* order. The species of the *Artemisietea vulgaris* class accompany this community.

Table 1. Floristic structure of the plant communities with *Solidago canadensis* L.

	Relevé number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
	Area [m ²]	100	100	100	100	50	44	100	100	100	60	100	100	100	100	100	100	100	100	100	60	100	100	100	100	100	100	100	100	
	Herb cover [%]	98	90	100	100	75	80	90	50	100	85	90	90	100	100	100	60	100	80	35	65	100	95	100	95	95	95	95		
	Mosses and lichens cover [%]																													
Ch. Ass. <i>Kochietum arenariae</i> Fijalk. 1978		1	+	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	2	•	•	•	•	•	•	
<i>Bassia laniflora</i> (S.G. Gimel.) A.J. Scott																														
Ch., D. All. <i>Koelerion glaucae</i> Volk 1931																														
<i>Koeleria glauca</i> (Spreng.) DC.																														
<i>Puccinellium oreoselinum</i> (L.) Moench																														
<i>Solidago virgaurea</i> L.																														
Ch. Cl. <i>Koelerio-Corynephoreta canescensii</i> Klik et Novák 1941																														
<i>Cladonia</i> sp.																														
<i>Helichrysum arenarium</i> (L.) Moench		1	2																											
<i>Festuca ovina</i> L.																														
<i>Jasione montana</i> L.																														
<i>Scleranthus perennis</i> L.																														
<i>Thymus serpyllum</i> L.																														
<i>Trifolium arvense</i> L.																														
<i>Trifolium campestre</i> Schreb.																														
Ch. All. <i>Arrhenatherion elatioris</i> Luquet 1926																														
Ch. O. <i>Arrhenatheretalia elatioris</i> Tx. 1931																														
<i>Achillea millefolium</i> L.																														
<i>Campanula patula</i> L.																														
<i>Dactylis glomerata</i> L.																														
<i>Daucus carota</i> L.																														
<i>Knautia arvensis</i> (L.) Coultr.																														
<i>Pimpinella saxifraga</i> L.																														
<i>Taraxacum campyloides</i> G.E. Haglund																														
Ch., D. Ass. <i>Ranunculo-Allopecuretum geniculati</i> R. Tx. 1937																														
<i>Allopecurus geniculatus</i> L.																														
<i>Potentilla repens</i> L.																														
<i>Rorippa sylvestris</i> (L.) Besser																														
<i>Inula britannica</i> L.																														
Ch. All. <i>Potentillion unciniae</i> Tx. 1947																														
Ch. O. <i>Potentillo-Polygonetalia avicularis</i> Tx. 1947																														
<i>Agrostis stolonifera</i> L.																												3	5	•
<i>Carex hirta</i> L.																												+	+	3
<i>Potentilla anserina</i> L.																												+	+	•
<i>Rumex crispus</i> L.																												•	+	•
Ch. Cl. <i>Molinio-Arrhenatheretea</i> Tx. 1937																												2	3	+
<i>Festuca pratensis</i> Huds.																												•	•	•
<i>Festuca rubra</i> L.																												•	•	1

Relevé number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29
Area [m ²]	100	100	100	50	44	100	100	100	60	100	100	100	100	100	100	100	100	100	100	60	100	100	100	100	100	100	100	100	
Herb cover [%]	98	90	100	100	75	80	90	50	100	85	90	90	100	100	100	100	100	100	100	80	35	65	100	95	100	95	95	95	
Mosses and lichens cover [%]																													
<i>Chamaecytisus ruthenicus</i> (Fisch. ex Wol.) Klask.	•	+	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Frangula alnus</i> Mill.	•	•	•	•	•	•	•	•	+	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Jurinea cyanooides</i> (L.) Rchb.	•	•	•	•	+	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Pinus sylvestris</i> L.	•	•	•	•	•	•	•	•	+	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	+	
<i>Pleurozium shreberi</i> (Willd ex Brid) Mitt.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
Cl. <i>Carpinio-Fagetea sylvatica</i> Jakucs ex Passarge 1968	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Seriphularia nodosa</i> L.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
Cl. <i>Alnetea glutinosae</i> Br.-Bl. et Tx. ex Westhoff et al. 1946	•	•	1	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Humulus lupulus</i> L.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Salix pentandra</i> L.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
Other species																													
<i>Hypericum perforatum</i> L.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Populus nigra</i> L.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	
<i>Populus tremula</i> L.	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	

Dates and localities of relevés:

1 – 11.07.2020; the pine-forest terrace of the Snov River, the town of Snovsk (the Chernihiv region); 2 – 11.07.2020; the pine-forest terrace of the Snov River, the Zheyeda village (the Snovsk district, Chernihiv region); 3 – 11.07.2020; the pine-forest terrace of the Snov River, the Zhavid village, the Khrenovka village (the Snovsk district, Chernihiv region); 5 – 15.09.2020; the pine-forest terrace of the Stryzhen River, the city of Chernihiv; 6 – 13.09.2020; the Ripyk-Chernihiv loess “island”, the city of Chernihiv; 7 – 13.09.2020; the meadow-swamp complex “Zamglay, the Lovin village (the Ripyk district, Chernihiv region); 8 – 15.09.2020; the pine-forest terrace of the Snov River, the town of Snovsk (the Chernihiv region); 9 – 29.07.2020; the Ripyk-Chernihiv loess “island”, the Novoselyivka village (the Chernihiv district, Chernihiv region); 10 – 11.07.2020; the flood lands of the Snov River, the Yausevo cemetery, the city of Chernihiv; 11 – 13.09.2020; the Ripyk-Chernihiv loess “island”, the Mykhailo-Kotsibynsky loess “island”, the Zhukotki village (the Chernihiv district, Chernihiv region); 15 – 20.08.2019; the meadow and swamp complex “Zamglay, the Lovin village (the Chernihiv district, Chernihiv region); 17 – 28.07.2019; the Berezna-Mena-Sosnytsia loess “island”, the town of Berezna (the Mena district, Chernihiv region); 19 – 15.09.2020; the pine-forest terrace of the Stryzhen River, the city of Chernihiv; 20 – 11.07.2020; the pine-forest terrace of the Snov River, the Zagrebelska Sloboda village (the Snovsk district, Chernihiv region); 21 – 15.09.2020; the pine-forest terrace of the Kryukova River, the Cheretske village (the Horodnya district, Chernihiv region); 22 – 29.07.2020; the flood lands of the Snov River, the Pishchanka village (the Chernihiv district, Chernihiv region); 23 – 11.07.2020; the pine-forest terrace of the Snov River, the Molozhava village (the Horodnya district, Chernihiv region); 24 – 06.09.2020; the Ripyk-Chernihiv loess “island”, the Novoselyivka village (the Chernihiv district, Chernihiv region); 25 – 11.07.2020; the flood lands of the Kryukova River, the Molozhava village (the Horodnya district, Chernihiv region); 27 – 13.09.2020; the Ripyk-Chernihiv loess “island”, the Yatsevo tract, the city of Chernihiv; 28 – 15.09.2020; the pine-forest terrace of the Stryzhen River, the city of Chernihiv;

Syntax:

1 – association *Berteroëtum incanæ* Siss. et Tideman in Siss. 1950; 2 – community: *Solidago canadensis* L. [Onopordion acanthii Br.-Bl. et al. 1936]; 3 – initial community: *Solidago canadensis* L. [Onopordion acanthii Br.-Bl. et al. 1936 + Epilobieae angustifoliae Tx. et Preisig ex von Rochow 1951]; 4 – initial community: *Solidago canadensis* L. [Onopordion acanthii Br.-Bl. et al. 1936+ Koeleria glaucae Volk 1931]; 5 – initial community: *Solidago canadensis* L. – *Calamagrostis epigejos* (L.) Roth [Onopordion acanthii Br.-Bl. et al. 1936 + Koelerio-Corynephoretea canescensitatis Klika in Klka et Novák 1941]; 7 – initial community: *Solidago canadensis* L. – *Calamagrostis epigejos* (L.) Roth [Onopordion acanthii Br.-Bl. et al. 1936 + Molinio-Arrhenatheretea Tx. 1937]; 8 – initial community: *Solidago canadensis* L. – *Elytrigia repens* (L.) Nevski [Onopordion acanthii Br.-Bl. et al. 1936 + Molinio-Arrhenatheretea Tx. 1937]; 11 – initial community: *Solidago canadensis* L. – *Convolvulo arvensis-Agropyrium repens* (L.) Nevski [Convolvulo arvensis-Agropyrium repens (L.) Nevski [Onopordion acanthii Br.-Bl. et al. 1936 + Molinio-Arrhenatheretea Tx. 1937]]; 12 – association *Koelerio-Astragalum arenarium* (L.) Roth [Koelerio-Astragalum arenarium (L.) Roth [Onopordion acanthii Br.-Bl. et al. 1936 + Molinio-Arrhenatheretea Tx. 1937]]; 13 – association *Kochietum arenariae* Fjalk. 1978; 14 – initial community: *Solidago canadensis* L. – *Erigeron annuus* (L.) Pers. [*Koelerio-Corynephoretea canescensitatis* Klka in Klka et Novák 1941 + Molinio-Arrhenatheretea Tx. 1937]; 15 – initial community: *Carex ryzina* Blytt ex Lindblom [*Koelerio-Corynephoretea canescensitatis* Klka in Klka et Novák 1941 + Molinio-Arrhenatheretea Tx. 1937]; 16 – initial community: *Festuca pratensis* Huds. – *Ambrosia artemisiifolia* L. [*Arrenatheretalia elatioris* Tx. 1931]; 17 – association *Ranunculo-Alpecuretea geniculati* R. Tx. 1937; 18 – initial community: *Solidago canadensis* L. – *Poa pratensis* L. [*Molinio-Arrhenatheretea* Tx. 1937].

■ – dominant species of the initial community.

Similar communities with *S. canadensis* participation, investigated in Kursk, Bryansk, and Minsk, were described by Arepeeva and Kulikova (2017). The distinctive feature of their relevés is the participation of the species, diagnosing the *Agropyretalia intermedio-repentis* order.

There is also a number of initial communities with characteristic species of the *Onopordetalia acanthii* order, where *S. canadensis* has a projective coverage from 20 to 98%, but in combination with characteristic species of other high-ranking taxa. Such species as *Equisetum arvense* (relevé 12), *Calamagrostis epigejos* (relevés 9, 11, 13), *Agrostis capillaris* (relevé 10), *Artemisia scoparia* (relevé 19) and *Elytrigia repens* (relevés 14–18) co-dominate in the initial communities. These communities are mostly the transformed meadow phytocenoses (relevés 7–11) and less often – the communities of pine-forest sands (relevés 5–6). A process of meadow ecosystems ruderalization due to invasion of *S. canadensis* is shown in the relevés 12–19. This process is especially manifested on the loess “islands” (relevés 15–18), where a gradual replacement of meadows with semiruderal grasslands and herblands of the *Convolvulo arvensis-Agropyron repens* alliance could be observed. The community, described in the relevé 24, is an example of the initial invasion of *S. canadensis* into the meadow phytocenosis, which had already been transformed by *Ambrosia artemisiifolia* – the species, which is widely represented in the Chernihiv-Ripky loess island phytocenoses.

The communities of the *Koelerio-Corynephoretea canescens* class and the *Koelerion glaucae* alliance were described in the pine-forest terraces of the Snov (near Zagrebelsna Sloboda village) and the Kryukova (near Cherepetske village) rivers. The vegetation of the *Koelerio-Corynephoretea canescens* class is widespread in a temperate zone of Europe. This sand vegetation is most abundant in northwestern Europe and in the lowlands of northern Germany and Poland. The pioneer communities are widespread under the conditions of xerophytization and anthropogenic transformation in the Chernihiv Polesie. The community (relevé 20) of the *Koelerio-Astragaletum arenarii* association is diagnosed by *Astragalus arenarius*. This species, which is included into the Red Book of the Ukraine, is rare in the Chernihiv Polesie. The *Kochietum arenariae* association, diagnosed by *Bassia laniflora*, is represented in the relevé 21. We noted earlier that the communities of synanthropic psammophytes species more resistant to anthropogenic influence, in particular *Kochietum arenariae*, are formed under urban conditions (Lukash and Danko, 2020).

Participation of *Solidago canadensis* in the psammophytic communities (relevés 20–21) of the *Koelerio-Corynephoretea canescens* class is not significant. Characteristic species of the *Vaccinio-Piceetea* class (*Jurinea cyanoides*, *Pinus sylvestris*, *Pleurozium shreberi*) also take part in the phytocenosis formation. Characteristic species of the *Koelerion glaucae* alliance (such as *Koeleria glauca*, *Peucedanum oreoselinum*, *Solidago virgaurea*) and the *Koelerio-Corynephoretea canescens* class (*Cladonia* sp., *Festuca ovina*, *Scleranthus perennis*, *Thymus serpyllum*) are represented in these two communities.

There are also transformed phytocenoses, possessing some features of the *Koelerio-Corynephoretea canescens* class at different succession stages. The relevé 22 (*Solidago canadensis-Erigeron annuus* initial community) illustrates the xerophytization process of the meadow floodplain phytocenoses grassland under conditions of the *S. canadensis* invasion. The initial community is formed by the dominant species: *Solidago canadensis* (with projective coverage 30%) and *Erigeron annuus* (25%). As part of these communities, there is a number of characteristic species of the *Koelerio-Corynephoretea canescens* (*Helichrysum arenarium*, *Festuca ovina*, *Jasione montana*, *Thymus serpyllum*) and *Molinio-Arrhenatheretea* (*Phleum pratense*, *Plantago lanceolata*, *Ranunculus acris*, *Rhinanthus minor*, *Rumex acetosa*, *Vicia cracca*) classes.

In the relevé 23, collected in the meadow and swamp complex Zamglay near Lovin village, characteristic species of the *Koelerio-Corynephoretea canescens* class (such as *Helichrysum arenarium*, *Festuca ovina*, *Thymus serpyllum*, *Trifolium arvense*, *Trifolium campestre*) and of the *Molinio-Arrhenatheretea* class (*Festuca pratensis*, *Plantago lanceolata*, *Rumex acetosa*, *Vicia cracca*) were identified. The *Carex rhizina* initial community (*Koelerio-Corynephoretea canescens* + *Molinio-Arrhenatheretea*) is formed by the monodominant species *Carex rhizina* with its projective coverage 90%. We think that the *Carex rhizina* initial community is a stage of overgrowth and xerophytization of the meadow-swamp complex Zamglay in the next succession series: marsh vegetation → semiruderal phytocenoses of nutrient-demanding grasses on anthropogenic transformed soils → meadow ruderal herblands with *Solidago canadensis* + other species → meadow ruderal herblands with monodomination of *Solidago canadensis* → xerophytic community.

All initial communities should be treated as successional phases of different phytocenoses under the impact of *Solidago canadensis* invasion. The *S. canadensis* spreading leads to the unification and loss of the phytocenoses traits. The relevés 8–10 represent the initial stages of successional changes under the influence of the *S. canadensis* invasion into the floodplains of small rivers, when phytocenosis originality still remains. Instead, on the loess islands this process is accelerated by the anthropogenic pressure on ecosystems and has irreversible consequences. For example, in the plant community *Poa pratensis-Festuca rubra*, *Solidago canadensis* firstly displaces *Festuca rubra* (relevé 29) and then forms the entire phytocenosis (relevés 27, 28).

CONCLUSIONS

The *Solidago canadensis* habitats in the Chernihiv Polesie are partially or completely ruderalized meadow phytocenoses of the river floodplains. Meadows with *S. canadensis* represent different stages of succession with a significant participation of characteristic species of the *Artemisieta vulgaris* class. Significant participation of the

Papaveretea rhoeidis class monocarpics testifies to the leading role of anthropogenic influence in the processes of xerophytization and unification of the meadow phytocenoses due to the *S. canadensis* invasion. Anthropogenic pressure on ecosystems accelerates the invasion of *S. canadensis* into phytocenoses and also leads to irreversible consequences. The *Berteroëtum incanae* association ruderal communities are the most common phytocenoses with *S. canadensis*. The *S. canadensis* invasion on the loess islands leads to a formation of semi-ruderal grasslands and herblands of the *Convolvulo arvensis–Agropyron repens* alliance. *S. canadensis* also occurs on pine terraces, in particular in the *Koelerio–Corynephoretea* class communities, which are mostly successive stages of transformation of the psammophytic herbal phytocenoses.

Thus, general trends in the distribution of communities with *Solidago canadensis* in the meadows and sands of the region are:

- Annual and biennial species would be well represented as a sign of recent disturbance of ecosystems.
- Floristic composition shows such tendencies in the development of plant communities with *S. canadensis* participation as xerophytization, ruderalization and species unification of the natural phytocenoses (of both psammophytic and meadow types).

Taking into consideration the peculiarities of the development cycle of *S. canadensis*, in order to prevent phytovinvasion, the authors consider it appropriate to carry out measures of plants mowing in places of their concentration at the beginning of mass flowering. Phytocenoses with *S. canadensis* need further attention as a possible source of late summer and autumn honey harvest.

REFERENCES

- Abhilasha, D., Quintana, N., Vivanco, J., Joshi, J., 2008. Do allelopathic compounds in invasive *Solidago canadensis* s.l. restrain the native European flora? *Journal of Ecology* 96, 993–1001.
- Arepieva, L.A., Kulikova, E.Ya., 2017. Communities with *Solidago canadensis* and *S. gigantea* in the cities of Kursk, Bryansk and Minsk. *Plant diversity* 3 (11), 38–43 (in Russian).
- Bielecka, A., Borkowska, L., Królik, E., 2020. Environmental changes caused by the clonal invasive plant *Solidago canadensis*. *Annales Botanici Fennici. Finnish Zoological and Botanical Publishing Board* 57 (1–3), 33–48.
- Burda, R.I., Pashkevich, N.A., Boyko, G.V., Fitsaylo, T.V., 2015. Alien species of natural flora of the Forest-Steppe and Steppe. *Scientific Opinion of the National Academy of Sciences of Ukraine*, Kyiv, 120 pp. (in Ukrainian).
- Daineka, M., Timofeev, S., 2018. Development of invasive species Canadian goldenrod (*Solidago canadensis* L.) in Vetka and Chech-
- ersk districts of Gomel region. *Bulletin of Science and Practice* 4 (4), 12–19 (in Russian).
- Dassonville, N., Vanderhoeven, S., Vanparys, V., Hayez, M., Gruber, W., Meerts, P., 2008. Impacts of alien invasive plants on soil nutrients are correlated with initial site conditions in NW Europe. *Oecologia* 157, 131–140.
- Dubovik, D.V., Skuratovich, A.N., Miller, D., Spiridovich, E.V., Gorbunov, Yu.N., Vinogradova, Yu.K., 2019. The invasiveness of *Solidago canadensis* in the Sanctuary «Prilepsky» (Belarus). *Nature Conservation Research* 4 (2), 48–56.
- Gusev, A.P., 2015. Impact of invasion of Canadian goldenrod (*Solidago canadensis* L.) on restorative succession in abandoned lands (southeast of Belarus). *Russian Journal of Biological Invasions* 6 (2), 74–77 (in Russian).
- Gusev, A.P., Shpileuskaya, N.S., 2016. Invasion of Canadian Goldenrod (*Solidago canadensis* L.) in a technogenic landscape (on example an open-cast mine on sand). *Bulletin of Polesie State University. Natural Sciences Series* 2, 3–7 (in Russian).
- Hennekens, S.M., Schaminée, J.H.J., 2001. Turboveg, a comprehensive database management system for vegetation data. *Journal of Vegetation Science* 12, 589–591.
- Lavrenko, E.M., Korchagin, A.G., 1976. Field geobotany. The structure of plant communities 5. Nauka, Leningrad, 320 pp. (in Russian).
- Lukash, O., Danko, H., 2020. The vegetation of sands in the Chernihiv city (Ukraine). *Studia Quaternaria* 37 (1), 31–44.
- Lukash, O., Yakovenko, O., Miroshnyk, I., 2018. The mechanical degradation of land surface and the present state of the loess “islands” plant cover of Chernihiv Polesie (Ukraine). *Ecological Questions* 29 (4), 23–34.
- Marynich, A.M., Parkhomenko, H.O., Petrenko, O.M., Shyshenko, P.H., 2003. Improved scheme of physical and geographical zoning of Ukraine. *Ukrainian Geographic Journal* 1 (41), 21–32 (in Ukrainian).
- Matuszkiewicz, W., 2019. Guide to the determination of Polish plant communities (Przewodnik do oznaczania zbiorowisk roślinnych Polski). Wydawnictwo Naukowe PWN, Warszawa, 404 pp. (in Polish).
- Mucina, L., Brandes, D., 1985. Communities of *Berteroë incana* in Europe and their geographical differentiation. *Vegetatio* 59, 125–136.
- Mucina, L., Bütlmann, H., Dierßen, K., Theurillat, J.-P., Raus, T., Čarní, A., Šumberová, K., Willner, W., Dengler, J., García, R.G., Chytrý, M., Hájek, M., Di Pietro, R., Iakushenko, D., Pallas, J., Daniëls, F.J.A., Bergmeier, E., Guerra, A.S., Ermakov, N., Valachovič, M., Schaminée, J. H.J., Lysenko, T., Didukh, Y.P., Pignatti, S., Rodwell, J.S., Capelo, J., Weber, H.E., Solomeshch, A., Dimitopoulos, P., Aguiar, C., Hennekens, S.M., Tichý, L., 2016. Vegetation of Europe: hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Applied Vegetation Science* 19 (S1), 3–264.
- Solomakha, V.A., Kostylov, O.V., Sheliaj-Sosonko, Yu.R., 1992. Synanthropic vegetation of Ukraine. Naukova dumka, Kyiv, 250 pp. (in Ukrainian).
- Stefanic, E., Puskadija, Z., Stefanic, I., Bubalo, D., 2003. Goldenrod: A valuable plant for beekeeping in north-eastern Croatia. *Bee World* 84, 88–92.
- Tichý, L., 2002. JUICE, software for vegetation classification, *Journal of Vegetation Science* 13, 451–453.