

The first trigonotarbid arachnid from Ukraine

JASON A. DUNLOP¹ and VITALY S. DERNOV²

¹ *Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Invalidenstrasse 43, D-10115 Berlin, Germany; e-mail: jason.dunlop@mfn.berlin*

² *Institute of Geological Sciences, National Academy of Sciences of Ukraine, Oles Honchar Street 55b, 01054 Kyiv, Ukraine; e-mail: vitalydernov@gmail.com*

ABSTRACT:

Dunlop, J.A. and Dernov, V.S. 2023. The first trigonotarbid arachnid from Ukraine. *Acta Geologica Polonica*, 73 (2), 181–187. Warszawa.

The extinct arachnid order Trigonotarbida Petrunkevitch, 1949 is reported here for the first time from Ukraine. The material consists of an opisthosoma preserved in ventral view from the upper Carboniferous (lower Moscovian; *Paralegoceras–Eowellerites* ammonoid zone) of the Gorlivka locality in the Donets Basin, eastern Ukraine. Formal assignment to a family or genus is difficult, but the preserved ventral anatomy is consistent with a member of the families Aphantomartidae Petrunkevitch, 1945, Kreischeriidae Haase, 1890 or Eophryniidae Karsch, 1882. It is noteworthy for expanding the known distribution of trigonotarbids in Europe and is only the second Palaeozoic arachnid to be formally described from Ukraine; the other being the carapace of a whip scorpion (Thelyphonida Latreille, 1804) from Lomovatka in the Luhansk Region, also in the Donets Basin.

Key words: Carboniferous; Moscovian; Arachnida; Trigonotarbida; Donets Basin; Ukraine.


INTRODUCTION

Trigonotarbids (Arachnida Lamarck, 1801) are an extinct order of spider-like arachnids. They belong to an arachnid clade named Pantetrapulmonata Shultz, 2007 sharing several features with spiders and their closest relatives including mouthparts shaped like a clasp-knife, a narrowing of the body between the prosoma and opisthosoma and the presence of two pairs of book lungs in the ground plan. For an overview of these animals see, e.g., Garwood and Dunlop (2010) or Jones *et al.* (2014). Seventy species of trigonotarbids are currently recognised (Dunlop *et al.* 2023), ranging in time from the late Silurian to the early Permian. Most species, and specimens, come from the upper Carboniferous/Pennsylvanian Coal Measures of Europe and North America with particularly strong concentrations in areas associated with coal mining. For a comprehensive list of localities and their stratigraphy see Dunlop and Rößler (2013, table 1 and fig. 6).

Here, we expand the known geographical range of Trigonotarbida by describing the first representative of this arachnid order from Ukraine. The specimen consists of an opisthosoma preserved in ventral view from the upper Carboniferous of the Donets Basin. It also represents only the second example of a Palaeozoic arachnid from Ukraine, the other being the unnamed carapace of a whip scorpion (Thelyphonida Latreille, 1804) described by Selden *et al.* (2014).

MATERIAL AND METHODS

The present fossil was discovered in the collections of the Institute of Geological Sciences of the National Academy of Sciences of Ukraine, Kyiv (IGS NASU). It was found in the core sample of borehole No. 118 drilled near the town of Gorlivka in the Donetsk Region, eastern Ukraine (Text-fig. 1) at a depth of 122.9 m. The single specimen (IGS



Text-fig. 1. Stratigraphic and geographic location of the trigonotarbid arachnid from Ukraine. A, B – Stratigraphic position of the fossil (orange star) in the Carboniferous succession of the Donets Basin. C–E – Maps indicating the general and detailed position of the locality. Abbreviations: Tour. – Tournaisian, Serpukhov. – Serpukhovian, Kasimov. – Kasimovian. Lithological column in B modified after Belokon (1963).


NASU-21/01) consists of an opisthosoma preserved in ventral view, plus a limb fragment (Text-fig. 2). The fossil-bearing rock is a dark gray coarse-grained carbonaceous and micaceous siltstone with clusters of carbonized plant debris. This siltstone lies 3.74 m above the I_4 coal layer in the middle part of the Almazna Formation.

The specimen was found in the 1940s or 1950s by Smirnova (forename unknown), a geologist for the ‘Uglerazvedka’ organization in the town of Artemovsk (now Bakhmut) in the Donetsk Region. The core sample, including the arachnid, was given to Dr. David E. Aisenverg (Institute of Geological Sciences, Academy of Sciences of the Ukrainian SSR, Kyiv), but for reasons unknown he did not complete

the study of this material and over time the specimen was misplaced. Only this year, a siltstone slab including the arachnid remains was found in the collection of Carboniferous brachiopods from the Donets Basin, stored in the Department of Palaeontology and Stratigraphy of the Palaeozoic sediments (IGS NASU, Kyiv).

GEOLOGICAL SETTING

The Almazna Formation (Text-fig. 1), also called C_2^6 or L, consists of a paralic succession of sandstones, siltstones, mudstones, coals and limestones. This formation lies on the sediments of the



Text-fig. 2. *Trigonotarbida incertae sedis*. A – IGS NASU-21/01 from the lower Moscovian (upper Carboniferous /Pennsylvanian) of Gorlivka in the Donets Basin, Ukraine. B – Interpretative drawing, opisthosomal segments numbered. Abbreviations: ap – anterior projection of sternites; gr – granular ornament; mb – marginal border; py – pygidium; sp? – possible marginal spines; IV – leg IV fragment (femur?). Scale bar equals 5 mm.

Kamenskaya Formation (also called C_2^5 or K) and is overlain by rocks of the Gorlivka Formation (also called C_2^7 or M). The limestone layer L_1 is a basal layer of the Almazna Formation; the upper boundary of this formation is at the base of the M_1 limestone layer. The thickness of this formation varies from 180 m in the NW part of the Donets Basin to 610 m in the SE part of the Donets Basin (Aisenverg *et al.* 1963; Dunayeva 1969; Nemyrovska and Yefimenko 2013). Near the town of Gorlivka its thickness is 270 m (Feofilova and Levenshtein 1963).

Alluvial sediments (42.8% of the section thickness), as well as sediments of peat and clastic swamps (10.8%), lacustrine (7.4%), lagoonal (26.0%), and shallow marine (13.0%) rocks are present in the section of the Almazna Formation (Feofilova and Levenshtein 1963, table 2). Alluvial sandstones are widespread in the lower part of this formation, in the interval between the L_1 and L_5 limestone layers. The upper part of this formation is dominated by marine

and lagoonal deposits, and among the sandstones, a significant portion belongs to deltaic sediments (Feofilova and Levenshtein 1963).

Coal-bearing deposits of the Moscovian strata in the Donets Basin were accumulated mainly in a large alluvial-deltaic plain, which was flooded periodically by the waters of warm epicontinental seas (Feofilova and Levenshtein 1963; Kozitskaya and Schegolev 1993). The Almazna Formation corresponds to the Mar'yivkian Horizon (upper half of the Lozovian Regional Stage) in the Regional stratigraphic scheme of the Dnipro-Donets Depression (Poletaev *et al.* 2011; Nemyrovska and Yefimenko 2013). The absolute age of the claystones (tonsteins in the original paper) from the I_3 coal bed, which lies c. 40 m below the I_4 coal bed, is 312.01 ± 0.08 and 312.18 ± 0.07 Ma (Davydov *et al.* 2010).

Sediments of the Almazna Formation contain conodonts of two zones: the '*Streptognathodus*' *transitivus*–*Neognathodus atokaensis* Zone (stratigraphic interval between the K_6 and L_5 limestone layers) and the younger *Idiognathodus izwaricus* Zone (L_5 – M_1 limestone layers) (Fohrer *et al.* 2007; Nemyrovska 2017). According to conodont studies (Nemyrovska 2017, text-fig. 2), the Mar'yivkian Horizon of the Donets Basin corresponds to the Kashirian Regional Stage of the stratotype of the Moscovian Global Stage (Moscow Syncline, Russia).

The Almazna Formation also contains remains of typical Bolsovian (= Westphalian C) terrestrial plants such as *Paripteris linguaeifolia* (Bertrand) Laveine, 1967, *Eusphenopteris striata* (Gothan) Novik, 1952, *Mariopteris nervosa* (Brongniart) Zeiller, 1879, *Macroneuropteris scheuchzeri* (Hoffmann) Cleal, Shute and Zoderow, 1990, *Laveineopteris rarineris* (Bunbury) Cleal, Shute and Zoderow, 1990, *Sphenophyllum emarginatum* Brongniart, 1822, etc. (Novik 1952, 1974; Fissunencko 1991, 2000; Boyarina 2016), as well as the nonmarine bivalves *Anthraconaia stobbsi* (Dix and Trueman, 1931), *A. oblonga* (Wright, 1929), *A. aff. pulchra* Hind, 1895, *Naiadites daviesi* (Dix and Trueman, 1931), *N. obliquus* Dix and Trueman, 1931, etc. (Sergeyeva 1981, 1984) and the ammonoids *Wiedeyoceras cf. cambriense* (Bisat, 1930); see also Derinov (in press). Additionally, Popov (1979) reported the Bolsovian ammonoid species *Wiedeyoceras cambriense* and *Politoceras cf. politum* (Shumard, 1858) from the lower part of the Gorlivka Formation. Other marine and terrestrial biota are also present in the Almazna Formation, including calcareous algae, microspores, foraminiferans, worms, corals, bryozoans, brachiopods, scaphopods, gastropods, horseshoe crabs, barnacles and ostracodes.

SYSTEMATIC PALEONTOLOGY

Order Trigonotarbida Petrunkevitch, 1949

Family *incertae sedis*

(Text-fig. 2)

MATERIAL: IGS NASU-21/01 from the lower Moscovian (upper Carboniferous /Pennsylvanian) of the Donets Basin, eastern Ukraine. Carbonaceous siltstones (Almazna Formation, 3.74 m above the l_4 coal layer) at a depth of 122.9 m of borehole No. 118 drilled near the town of Gorlivka (Donetsk Region, Ukraine); leg. Smirnova.

DESCRIPTION: Posterior half of a rounded to oval opisthosoma in ventral view (Text-fig. 2). Preserved length c. 12 mm; maximum width 14.8 mm. Sternites 5–10 visible, plus the two circular segments forming the pygidium. Sternites strongly procurved, all with a slight anterior projection close to the midline giving them a distinctive outline. Hints of light granulation towards the lateral margins, but sternites otherwise smooth. Opisthosoma with narrow (c. 0.2 mm) marginal border. Posterior margin broken in places; weak hints of posterior spines preserved, but their precise presence and form is equivocal. Pygidium forms two small concentric rings, diameter 1.6 mm, positioned towards anterior end of sternite 10. Limb fragment to the right of opisthosoma probably represents the femur of leg IV.

DISCUSSION

The incomplete nature of the new fossil makes it difficult to assign taxonomically, but it is undoubtedly a member of the extinct arachnid order Trigonotarbida as it has the characteristic ventral anatomy of this group such as the small, rounded segments forming a discrete pygidium on the ventral surface of the opisthosoma (py in Text-fig. 2). Seven trigonotarbid families are currently known from the Carboniferous and several can be excluded from consideration based on their morphology. Anthracomartidae Haase, 1890 usually have sternites which come to a sharp angle on the midline (Garwood and Dunlop 2011, text-figs 1–3) and additionally have a broad sulcus defining a much wider ventral lateral margin. Anthracosironidae Pocock, 1903 have a more elongate, somewhat pear-shaped opisthosoma (Pocock 1903, 1911; Petrunkevitch 1949) and thus presents a quite different habitus. Lissomartidae Dunlop, 1995 and Trigonotarbidae have procurved sternites with

smooth borders (Pocock 1911; Dunlop 1995; Jones *et al.* 2014), i.e., they do not show the anterior projection on the midline (ap in Text-fig. 2) preserved in the present fossil.

The best matches to the present specimen thus come from the families Aphantomartidae Petrunkevitch, 1945, Kreischeriidae Haase, 1890 and Eophrynidae Karsch, 1882. In species where the ventral opisthosoma is known, the sternites can also show a projection on the midline. Examples include the Devonian aphantomartid *Alkenia mirabilis* Størmer, 1970 and the Permian species *Aphantomartus ifeldicus* (Scharf, 1924) as figured by Poschmann and Dunlop (2010, fig. 1d) and Rößler *et al.* (2003, fig. 1), respectively. However, if the observation of posterior spines on the opisthosoma of the new fossil is correct (sp? in Text-fig. 2) this would tend to exclude aphantomartids which do not show such marginal spination and also tend to have a more obviously oval opisthosoma as opposed to a broader and more rounded one; see also figures in Rößler (1998).

Among the kreischeriids this ventral morphology with midline projections is seen in *Kreischeria wiedeii* Geinitz, 1882 and *Pseudokreischeria pococki* (Gill, 1924); see especially Rößler and Dunlop (1997, fig. 4) and Dunlop (1998, fig. 4), respectively. This morphology also appears to be present in the kreischeriid *Gondwanarachne argentinensis* Pinto and Hünicken, 1980, the only trigonotarbid currently known from the Southern Hemisphere. Interestingly, the preserved ventral morphology based on Pinto and Hünicken (1980, fig. 3) is quite similar to the new fossil with a narrow marginal border, longitudinal lines towards the margins at an angle and hints of marginal granulation. A median projection of the sternites is also seen, albeit perhaps less strongly, in eophrynids such as *Eophrynus prestvicii* (Buckland, 1837), *Stenotrogulus salmii* (Stur, 1877) and *Pleophrynus verrucosus* (Pocock, 1911). Figured examples for the respective species include Dunlop and Garwood (2014, fig. 2B), Příbyl (1958, figs 1, 2) for a synonym of *S. slammii*, and Dunlop (1994, figs 1b, 2a). Aphantomartidae, Kreischeriidae and Eophrynidae are probably closely related (Jones *et al.* 2014) and are best differentiated in dorsal view based on patterns of tuberculation on the opisthosomal tergites. Since this character is not preserved here, we cannot unequivocally resolve which of these families the new Ukrainian fossil belongs to.

It is nevertheless important for documenting the presence of Trigonotarbida at a locality further east than some of the main concentrations of records in, e.g., the Coal Measures of Silesia in Poland and Bohemia in the Czech Republic. Together with the

whip scorpion described by Selden *et al.* (2014) they suggest that the Donets Basin has considerable potential to provide useful comparative data about Carboniferous arachnid faunas across a wider region of the original paleocontinent of Laurasia. We should also reiterate that these two Carboniferous fossils are the oldest arachnids known from Ukraine. The only other source of fossil arachnids from Ukraine is the Paleogene Rovno amber (e.g., Wunderlich 2004; Perkovsky *et al.* 2007; Dunlop *et al.* 2019), which has yielded a rich assemblage of spiders, scorpions, harvestmen and mites; albeit mostly in modern families and/or genera.

Acknowledgements

We thank Dr. Vladyslav Poletaev (IGS NASU, Kyiv) for his help with the historical collections, and Markus Poschmann and an anonymous reviewer for helpful comments on an earlier draft of the typescript.

REFERENCES

- Aisenverg, D.E., Brazhnikova, N.E., Novik, K.O., Rotai, A.P. and Shulga, P.L. 1963. Carboniferous stratigraphy of the Donets Basin, 183 pp. Publishing House of the Academy of Sciences of the Ukrainian SSR; Kyiv. [In Russian]
- Belokon, V.G. 1963. Central coal-bearing district. In: Kuznetsov, I.A. (Ed.), *Geology of coal and oil shale deposits of the European part of the USSR*, 670–687. Gostoptekhizdat; Moscow. [In Russian]
- Bisat, W.S. 1930. On the goniatite and nautiloid fauna of the Middle Coal Measures of England and Wales. *Summary of progress of the Geological Survey of Great Britain and the Museum of Practical Geology for the Year 1929*, **3**, 75–90.
- Boyarina, N.I. 2016. Middle and Upper Carboniferous (Pennsylvanian) megafloral zones of the Donets Basin. *Geologichnyy zhurnal*, **354**, 21–35. [In Russian]
- Brongniart A. 1822. Sur la classification et la distribution des végétaux fossiles. *Mémoires du Muséum d'histoire naturelle*, **8**, 203–240.
- Buckland, W. 1837. The Bridgewater treatises on the power, wisdom and goodness of God as manifested in the creation. Treatise IV. *Geology and mineralogy with reference to natural theology*. 2nd Edition, 747 pp. William Pickering; London.
- Cleal, C.J., Shute, C.H. and Zодrow, E.L. 1990. A revised taxonomy for Palaeozoic neuropterid foliage. *Taxon*, **39**, 486–492.
- Davydov, V.I., Crowley, J.L., Schmitz, M.D. and Poletaev, V.I. 2010. High-precision U-Pb zircon age calibration of the global Carboniferous time scale and Milankovitch band cyclicity in the Donets Basin, eastern Ukraine. *Geochemistry, Geophysics, Geosystems*, **11**, 1–22.
- Dernov, V.S. in press. Ammonoid biostratigraphy of the Bashkirian (Pennsylvanian) in the Donets Basin. *Collection of scientific works of the Institute of Geological Sciences of NAS of Ukraine*, **15**. [In Ukrainian]
- Dix, E. and Trueman, A.E. 1931. Some nonmarine lamellibranchs from the upper part of the Coal Measures. *Quarterly Journal of the Geological Society*, **87**, 180–211.
- Dunayeva, N.M. 1969. Open Donets Basin. In: Bondarchuk, V.G. (Ed.), *Stratigraphy of the Ukrainian SSR. Volume V. Carboniferous*, 21–48. Naukova Dumka; Kyiv. [In Ukrainian]
- Dunlop, J.A. 1994. The palaeobiology of the Writhlington trigonotarbid arachnid. *Proceedings of the Geologists' Association*, **105**, 287–296.
- Dunlop, J.A. 1995. Redescription of the Pennsylvanian trigonotarbid arachnid *Lissomartus Petrunkevitch* 1949 from Mazon Creek, Illinois. *Journal of Arachnology*, **23**, 118–124.
- Dunlop, J.A. 1998. A redescription of the trigonotarbid arachnid *Pseudokreischeria pococki* (Gill, 1924). *Bulletin of the British arachnological Society*, **11**, 49–53.
- Dunlop, J.A. and Garwood, R.J. 2014. Tomographic reconstruction of the exceptionally preserved trigonotarbid arachnid *Eophrynus prestvicii*. *Acta Palaeontologica Polonica*, **59**, 443–454.
- Dunlop, J.A., Marusik, Y. and Vlaskin, A.P. 2019. Comparing arachnids in Rovno amber with the Baltic and Bitterfeld deposits. *Paleontological Journal*, **53**, 1074–1083.
- Dunlop, J.A., Penney, D. and Jekel, D. 2023. A summary list of fossil spiders and their relatives. In: *World Spider Catalog*. Natural History Museum Bern, online at <http://wsc.nmbe.ch>, version 23.5, accessed on 01.03.2023.
- Dunlop, J.A. and Rößler, R. 2013. The youngest trigonotarbid *Permotarbus schuberti* n. gen., n. sp. from the Permian Petrified Forest of Chemnitz in Germany. *Fossil Record*, **16**, 229–243.
- Feofilova, A.P. and Levenshtein, M.L. 1963. Features of the sedimentation and coal accumulation in the Early and Middle Carboniferous of the Donets Basin, 175 pp. Publishing House of the Academy of Sciences of the USSR; Moscow. [In Russian]
- Fissunenکو, O.P. 1991. Zonal phytostratigraphical scale of the Lower and Middle Carboniferous of the Donets Basin. *Geologichnyy zhurnal*, **258** (3), 55–64. [In Russian]
- Fissunenکو, O.P. 2000. On the problem of the Moscovian Stage, 66 pp. Publishing House of Luhansk University; Luhansk. [In Russian]
- Fohrer, B., Nemyrovska, T.I., Samankassou, E. and Ueno, K. 2007. The Pennsylvanian (Moscovian) Izvarino section, Donets Basin, Ukraine: a multidisciplinary study on microfacies, biostratigraphy (conodonts, foraminifers, and ostracodes), and paleoecology. *Journal of Paleontology*, **81** (5), 1–85.

- Garwood, R.J. and Dunlop, J.A. 2010. Fossils explained 58. Trigonotarbids. *Geology Today*, **26**, 34–37.
- Garwood, R.J. and Dunlop, J.A. 2011. Morphology and systematics of Anthracomartidae (Arachnida: Trigonotarbida). *Palaeontology*, **54**, 145–161.
- Geinitz, H.B. 1882. *Kreischeria wiedeii*, ein Pseudoskorpion aus der Steinkohlenformation von Zwickau. *Zeitschrift der Deutschen geologischen Gesellschaft*, **34**, 238–242.
- Gill, E.L. 1924. Fossil arthropods from the Tyne Coalfield. *Geological Magazine*, **61**, 445–471.
- Haase, E. 1890. Beitrag zur Kenntniss der fossilen Arachniden. *Zeitschrift der Deutsche geologische Gesellschaft*, **1890**, 629–657.
- Hind, W. 1894–1896 A monograph on *Carbonicola*, *Anthracomya*, and *Naiadites*, 182 pp. Printed for the Palaeontographical Society; London.
- Jones, F.M., Dunlop, J.A., Friedman, M. and Garwood, R.J. 2014. *Trigonotarus johnsoni* Pocock, 1911, revealed by X-ray computed tomography, with a cladistic analysis of the extinct trigonotarbid arachnids. *Zoological Journal of the Linnean Society*, **172**, 49–70.
- Karsch, F. 1882. Ueber ein neues Spinnenthier aus der Schlesischen Steinkohle und die Arachniden überhaupt. *Zeitschrift der Deutschen geologischen Gesellschaft*, **34**, 556–561.
- Kozitskaya, R.I. and Schegolev, A.K. 1993. Late Carboniferous. In: Tsegelnyuk, P.D. (Ed.), Geological history of the territory of Ukraine, 147–154. IGS NASU; Kyiv. [In Russian]
- Lamarck, J.B.P.A. 1801. *Système des animaux sans vertèbres*, 432 pp. Deterville; Paris.
- Latreille, P.A. 1804. *Histoire naturelle, generale et particuliere, des Crustacés et des Insectes*, Vol. 7, 456 pp. Dufart; Paris.
- Laveine, J.-P. 1967. Contribution à l'étude de la flore du terrain houiller: les neuroptéridées du Nord de la France, 344 pp. Imprimerie Central du Nord; Lille.
- Nemyrovska, T.I. 2017. Late Mississippian–Middle Pennsylvanian conodont zonation of Ukraine. *Stratigraphy*, **14**, 299–217.
- Nemyrovska, T.I. and Yefimenko, V.I. 2013. Middle Carboniferous (Lower Pennsylvanian). In: Gozhik, P.F. (Ed.), Stratigraphy of the Upper Proterozoic and Phanerozoic of Ukraine. Volume 1. Stratigraphy of the Upper Proterozoic, Paleozoic and Mesozoic, 283–303. Logos; Kyiv. [In Ukrainian]
- Novik, E.O. 1952. Carboniferous flora of the European part of the USSR, 468 pp. Publishing House of the Academy of Sciences of the USSR; Moscow. [In Russian]
- Novik, E.O. 1974. Regularities of development of the Carboniferous flora of the south of the European part of the USSR, 140 pp. Naukova Dumka; Kyiv. [In Russian]
- Perkovsky, E.E., Rasnitsyn, A.P., Vlaskin, A.P. and Taraschuk, M.V. 2007. A comparative analysis of the Baltic and Rovno amber arthropod faunas: representative samples. *African Invertebrates*, **48**, 229–245.
- Petrunkévitch, A.I. 1945. Palaeozoic Arachnida. An inquiry into their evolutionary trends. *Scientific Papers, Illinois State Museum*, **3** (2), 1–76.
- Petrunkévitch, A.I. 1949. A study of Palaeozoic Arachnida. *Transactions of the Connecticut Academy of Arts and Sciences*, **37**, 69–315.
- Pinto, I.D. and Hünicken, M.A. 1980. *Gondwanarachne* a new genus of the order Trigonotarbida (Arachnida) from Argentina. *Boletín de la Academia Nacional de Ciencias Córdoba*, **53**, 307–315.
- Pocock, R.I. 1903. A new Carboniferous arachnid. *Geological Magazine*, **4** (10), 247–251.
- Pocock, R.I. 1911. A monograph of the terrestrial Carboniferous Arachnida of Great Britain. *Monographs of the Palaeontographical Society*, **64**, 1–84.
- Poletaev, V.I., Vdovenko, M.V., Shchogolev, O.K., Boyarina, N.I. and Makarov, I.A. 2011. Stratotypes of the Carboniferous and Lower Permian regional stratigraphic units of the Dnipro-Donets Depression, 236 pp. Logos; Kyiv. [In Ukrainian]
- Popov, A.V. 1979. Carboniferous ammonoids of the Donets Basin and their stratigraphic significance. *Trudy VSEGEI, n.s.*, **220**, 1–119. [In Russian]
- Poschmann, M. and Dunlop, J.A. 2010. Trigonotarbid arachnids from the Lower Devonian (Lower Emsian) of Alken an der Mosel (Rhineland-Palatinate, SW Germany). *Paläontologische Zeitschrift*, **84**, 467–484.
- Příbyl, A. 1958. Some new Carboniferous arachnids from the Ostrava-Karviná coal district. *Časopis pro Mineralogii a Geologii*, **3**, 425–434.
- Rößler, R. 1998. Arachniden-Neufunde im mitteleuropäischen Unterkarbon bis Perm – Beitrag zur Revision der Familie Aphantomartidae Petrunkevitch 1945 (Arachnida, Trigonotarbida). *Paläontologische Zeitschrift*, **72**, 67–88.
- Rößler, R. and Dunlop, J.A. 1997. Redescription of the largest trigonotarbid arachnid – *Kreischeria wiedeii* Geinitz 1882 from the Upper Carboniferous of Zwickau, Germany. *Paläontologische Zeitschrift*, **71**, 237–245.
- Rößler, R., Dunlop, J.A. and Schneider, J. 2003. A redescription of some poorly known Rotliegend arachnids from the Lower Permian (Asselian) of the Ilfeld and Thuringian Forest Basins. *Paläontologische Zeitschrift*, **77**, 417–427.
- Scharf, W. 1924. Beitrag zur Geologie des Steinkohlengebietes im Südharz. *Jahrbuch des Halleschen Verbands für die Erforschung der Mitteldeutschen Bodenschätze und ihrer Verwaltung*, **4**, 404–437.
- Selden, P.A., Shcherbakov, D.E., Dunlop, J.A. and Eskov, K.Y. 2014. Arachnids from the Carboniferous of Russia and Ukraine, and the Permian of Kazakhstan. *Paläontologische Zeitschrift*, **88**, 297–307.
- Sergeyeva, M.T. 1981. Bashkirian (Middle Carboniferous) bivalves of the Donets Basin and their stratigraphic significance. *Tectonics and stratigraphy*, **21**, 53–61. [In Russian]

- Sergeyeva, M.T. 1984. On the evolution of the Carboniferous nonmarine bivalves of the USSR. In: Vialov, O.S. (Ed.), Paleontology and stratigraphy of the Phanerozoic of Ukraine, 49–53. Naukova Dumka, Kyiv. [In Russian]
- Shultz, J.W. 2007. A phylogenetic analysis of the arachnid orders based on morphological characters. *Zoological Journal of the Linnean Society*, **150**, 221–265.
- Shumard, B.F. 1858. Description of new fossils from the Coal Measures of Missouri and Kansas. *Transactions of the St. Louis Academy of Science*, **1**, 198–227.
- Størmer, L. 1970. Arthropods from the Lower Devonian (Lower Emsian) of Alken an der Mosel, Germany. Part 1: Arachnida. *Senckenbergiana lethaea*, **51**, 335–369.
- Stur, D. 1877. Die Culm-Flora der Ostrauer und Waldenburger Schichten. *Abhandlung der königliche geologische Reichsanstalt*, **4**, 5.
- Wright, W. B. 1929. The zonal succession in the Coal Measures around Manchester. *Summary of progress of the Geological Survey of Great Britain and the Museum of Practical Geology for the Year 1928*, **2**, 36–55.
- Wunderlich, J. 2004. Fossil spiders (Araneae) in Early Tertiary amber from the Ukraine. In: Wunderlich, J. (Ed.), *Beiträge zur Araneologie*, **3**, 1821–1829.
- Zeiller, R. 1879. Vegetaux fossiles du terrain Houiller de la France. *Explication de la carte géologique de la France*, **4**, 1–185.

Manuscript submitted: 16th February 2023

Revised version accepted: 13th March 2023